arXiv:1603.08583v1 [math.CO] 28 Mar 2016

ON A CONJECTURE OF CIGLER

D. STANTON

Abstract. A conjecture of Cigler which realizes normalized q-Hermite polynomials as moments is verified.

1. Introduction

The q-Hermite polynomials $H_n(x|q)$ may be defined by

$$H_n(x|q) = \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix}_q e^{-i(n-2k)\theta}, \quad x = \cos\theta.$$

(We have used the standard notation for q-series given in [2] and [3].) In [4, Th. 4.1], four moment representations are given for four different normalizations of these polynomials

$$H_n(x|q), \quad \frac{H_n(x|q)}{(q;q)_n}, \quad \frac{H_n(x|q^2)}{(q;q)_n}, \quad \frac{H_n(x|q^2)}{(-q;q)_n}.$$

Four other moment representations are alluded to at the end of Section 4 of [4],

$$\frac{H_{2n}(x|q)}{(q^2;q^2)_n}, \quad \frac{H_{2n}(x|q)}{(q;q^2)_n}, \quad \frac{H_{2n+1}(x|q)}{(q^2;q^2)_n}, \quad \frac{H_{2n+1}(x|q)}{(q^3;q^2)_n}.$$

Cigler [1, Section 3.4] conjectured that the 6th and 8th of these moment sequences may be combined into a single moment sequence

$$\mu_n = P_n(a) = \frac{1}{(q;q^2)_{[(n+1)/2]}} \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix}_q a^k, \quad n \geq 0$$

for a specific set of orthogonal polynomials. In this short note we prove Cigler's conjecture.

2. The conjecture

First we define a set of monic orthogonal polynomials $s_n(x)$ via a three-term recurrence relation.

Definition 2.1. *Let $s_n(x)$ be monic polynomials in x defined by*

$$s_{n+1}(x) = (x - b_n)s_n(x) - \lambda_n s_{n-1}(x), \quad n \geq 0, \quad s_0(x) = 1, \ s_{-1}(x) = 0,$$

Date: March 30, 2016.
The author was supported by NSF grant DMS-1148634.

where

$$b_n = -\frac{(1-q)}{(1-q^{2n+1})(1-q^{2n-1})(1+a)}\times \left(a\frac{(1-q^{2n-1})(1-q^{n+1})(1-q^n)}{1-q} - q^n\left(\frac{1-q^{n-1}}{1-q}+q^{n+1}\frac{1-q^n}{1-q}\right)(1+a)^2\right),\ \textit{for } n\geq 0 \textit{ even},$$

$$b_n = \frac{(1-q)}{(1-q^{2n+1})(1-q^{2n-1})(1+a)}\times \left(a\frac{(1-q^{2n+1})(1-q^{n-1})(1-q^n)}{1-q} - q^{n+1}\left(\frac{1-q^n}{1-q}+q^{n-2}\frac{1-q^{n+1}}{1-q}\right)(1+a)^2\right),\ \textit{for } n\geq 1 \textit{ odd},$$

$$\lambda_n = \frac{q^n(1+a)^2(1-q^{n-1})(1-q^n)}{(1-q^{2n-1})^2},\ \textit{for } n\geq 1 \textit{ even},$$

$$\lambda_n = -\frac{(a+q^n)(a+q^{n-1})(1+aq^{n-1})(1+aq^n)}{(1+a)^2(1-q^{2n-1})^2},\ \textit{for } n\geq 1 \textit{ odd}.$$

Definition 2.2. *Let L be the linear functional on polynomials in x given by*

$$L(1)=1,\quad L(s_n(x))=0,\quad n\geq 1.$$

Cigler [1, Section 3.4] conjectured

Conjecture 2.3. *(Cigler)*

$$L(x^n)=P_n(a).$$

3. The proof

To prove Conjecture 2.3, we give another basis for polynomials which has an explicit L value.

Proposition 3.1. *Cigler's conjecture is equivalent to*

$$L\left(\prod_{i=0}^{n-1}(x^2-a^2q^{2i})\right)=\frac{(-a;q)_{2n}}{(q;q^2)_n},$$

$$L\left(x\prod_{i=0}^{n-1}(x^2-a^2q^{2i})\right)=\frac{(-a;q)_{2n+1}}{(q;q^2)_{n+1}}.$$

Proof. We show that $L(x^n)=P_n(a)$ implies the L values which are given in Proposition 3.1. Let $\epsilon=0$ for the first case and $\epsilon=1$ for the second case. Expanding by the q-binomial theorem we see that

$$L\left(x^\epsilon\prod_{i=0}^{n-1}(x^2-a^2q^{2i})\right)=\sum_{k=0}^{n}\begin{bmatrix}n\\k\end{bmatrix}_{q^2}(-1)^k a^{2k}q^{2\binom{k}{2}}L(x^{2(n-k)+\epsilon})$$

$$=\sum_{k=0}^{n}\begin{bmatrix}n\\k\end{bmatrix}_{q^2}(-1)^k a^{2k}q^{2\binom{k}{2}}\frac{1}{(q;q^2)_{n-k+\epsilon}}\sum_{s=0}^{2(n-k)+\epsilon}\begin{bmatrix}2(n-k)+\epsilon\\s\end{bmatrix}_q a^s$$

$$=(q^2;q^2)_n\sum_{p=0}^{2n+\epsilon}\frac{a^p}{(q;q)_{2n+\epsilon-p}}\sum_{k=0}^{[p/2]}\frac{(-1)^k q^{2\binom{k}{2}}}{(q^2;q^2)_k(q;q)_{p-2k}}$$

$$=\frac{1}{(q;q^2)_{n+\epsilon}}\sum_{p=0}^{2n+\epsilon}a^p\begin{bmatrix}2n+\epsilon\\p\end{bmatrix}_q q^{\binom{p}{2}}=\frac{(-a;q)_{2n+\epsilon}}{(q;q^2)_{n+\epsilon}}.$$

The k-sum was evaluated by a limiting case of the q-Vandermonde sum, and the p-sum evaluated by the q-binomial theorem, see [2, Appendix]. □

Because we know the values of $L(s_n(x))$, if we can expand $\prod_{i=0}^{n-1}(x^2-a^2q^{2i})$ in terms of $s_n(x)$ we can verify Proposition 3.1. The next Proposition accomplishes this task.

Proposition 3.2. *For any non-negative integer n,*

$$\prod_{i=0}^{n-1}(x^2-a^2q^{2i}) = \sum_{k=0}^{2n} a_k^{(n)} s_{2n-k}(x),$$

where

$$\begin{aligned} a_{2k}^{(n)} =& \frac{(-aq^{2n-1};1/q)_{2k}}{(q^{4n-2k-1};1/q^2)_k} \begin{bmatrix} n \\ k \end{bmatrix}_{q^2}, \\ a_{2k+1}^{(n)} =& (1+a)\frac{(-aq^{2n-1};1/q)_{2k}}{(q^{4n-2k-1};1/q^2)_{k+1}} \begin{bmatrix} n \\ k+1 \end{bmatrix}_{q^2} (1-q^{2(k+1)}). \end{aligned}$$

Proof. Use induction on n and the three-term recurrence relation in Definition 2.1 twice. We must verify that

$$\begin{aligned} a_k^{(n+1)} =& -a^2q^{2n}a_{k-2}^{(n)} + a_k^{(n)} \\ &+ (b_{2n-k+2}+b_{2n-k+1})a_{k-1}^{(n)} \\ &+ (\lambda_{2n-k+3}+b_{2n-k+2}^2+\lambda_{2n-k+2})a_{k-2}^{(n)} \\ &+ (b_{2n-k+3}\lambda_{2n-k+3}+\lambda_{2n-k+3}b_{2n-k+2})a_{k-3}^{(n)} \\ &+ \lambda_{2n-k+4}\lambda_{2n-k+3}a_{k-4}^{(n)}. \end{aligned}$$

Since each of the terms is explicitly known, this tedious computation may be done by hand or computer. □

Theorem 3.3. *Cigler's conjecture is true.*

Proof. Proposition 3.2 shows that the first part of Proposition 3.1 holds,

$$L\left(\prod_{i=0}^{n-1}(x^2-a^2q^{2i})\right) = a_{2n}^{(n)} = \frac{(-a;q)_{2n}}{(q;q^2)_n}.$$

For the second part of Proposition 3.1, we find the constant term in the expansion of $x\prod_{i=0}^{n-1}(x^2-a^2q^{2i})$. Using Definition 2.1 this is

$$\begin{aligned} L\left(x\prod_{i=0}^{n-1}(x^2-a^2q^{2i})\right) =& a_{2n}^{(n)}b_0 + a_{2n-1}^{(n)}\lambda_1 \\ =& \frac{(-a;q)_{2n+1}}{(q;q^2)_{n+1}}. \end{aligned}$$

□

The Hankel determinant of these moments [1, Conjecture 3.1 (3.4)] is thereby evaluated.

Corollary 3.4. *The Hankel determinant is*

$$det(P_{i+j}(a))_{0\leq i,j\leq n} = \prod_{i=1}^{n} \lambda_i^{n+1-i}.$$

where λ_i is given in Definition 2.1.

Additional sets of classical orthogonal polynomials as moments are given in [4], [5] and [6].

References

[1] J. Cigler, Elementary observations on Rogers-Szegö polynomials, arxiv preprint, 1602.07850v1.pdf, February, 2016.

[2] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition Cambridge University Press, Cambridge, 2004.

[3] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one variable*, Cambridge University Press, paperback edition, Cambridge, 2009.

[4] M. E. H. Ismail and D. Stanton, q-Integral and moment representations for q-orthogonal polynomials, *Can. J. Math.*, 54 (2002), p. 709-735.

[5] M. E. H. Ismail and D. Stanton, Classical orthogonal polynomials as moments, Canad. J. Math. 49 (1997), no. 3, p. 520-542.

[6] M. E. H. Ismail and D. Stanton, More orthogonal polynomials as moments, Mathematical essays in honor of Gian-Carlo Rota (Cambridge, MA, 1996), p. 377-396, Progr. Math., 161, Birkhäuser Boston, Boston, MA, 1998.

School of Mathematics, University of Minnesota, Minneapolis, MN 55455
E-mail address: stanton@math.umn.edu